

East @ 5838 E. Central Ave. 316-440-4595 fax 316-440-4596 South @ 7030 S. Broadway St. 316-558-5950 fax 316-558-5951

IMCWichita.com

Acetaminophen/Ibuprofen Dosage Chart

Tylenol/Advil/Motrin

ACETAMINOPHEN: Give every 6 hours as needed: do NOT exceed 4 doses in 24 hours.

WEIGHT	ELIXIR	CHEWABLE	JR. STRENGTH	REG. STRENGTH
pounds (lbs.)	I teaspoon = 160mg/5ml	I tablet = 80mg	I caplet = 160mg	I tablet = 325mg
6-11 lbs	I/4 teaspoon (1.25 ml)			
12-17 lbs	I/2 teaspoon (2.5 ml)			
18-23 lbs	3/4 teaspoon (3.75 ml)			
24-35 lbs	I teaspoon (5 ml)	2 tablets		
36-47 lbs	I I/2 teaspoons (7.5 ml)	3 tablets		
48-59 lbs	2 teaspoons (10 ml)	4 tablets	2 caplets	l tablet
60-71 lbs	2 1/2 teaspoons (12.5 ml)	5 tablets	2 I/2 caplets	I tablet
72-95 lbs	3 teaspoons (15 ml)	6 tablets	3 caplets	I I/2 tablet
96+ lbs			4 caplets	2 tablets

IBUPROFEN: Give every 6-8 hours as needed always with food; do NOT exceed 4 doses in 24 hours.

WEIGHT	INFANT DROPS	CHILDREN'S LIQUID	CHEWABLE TABS	TABLETS
pounds (lbs.)	40mg/Iml	I teaspoon = 100mg/5ml	I tablet = 100 mg	I tablet = 200mg
11-21 lbs	1/4 teaspoon (1.25ml)	I/2 teaspoon (2.5ml)	I/2 tablet	
22-32 lbs	1/2 teaspoon (2.5ml)	I teaspoon (5ml)	l tablet	
33-43 lbs		I I/2 teaspoons (7.5ml)	I I/2 tablets	
44-54 lbs		2 teaspoons (10ml)	2 tablets	l tablet
55-65 lbs		2 1/2 teaspoons (12.5ml)	2 I/2 tablets	
66-87 lbs	7	3 teaspoons (15ml)	3 tablets	I I/2 tablets
88+ lbs		4 teaspoons (20ml)	4 tablets	2 tablets